



**Connecticut
Public Health
Association**

Promoting Public Health in Connecticut Since 1916

**TESTIMONY OF CONNECTICUT PUBLIC HEALTH ASSOCIATION
REGARDING SB 16, AN ACT REQUIRING THE LABELING OF FOOD
PACKAGING THAT CONTAINS BISPHENOL-A**

COMMITTEE ON ENVIRONMENT

February 25, 2013

Kathi Traugh
President

Katharine Kranz Lewis
President Elect

Alison Gilcreast
Treasurer

Kimberly Ploszaj
Secretary

Tracey Scraba
Immediate Past-President

Board of Directors

- Elaine Abrams
- Ashika Brinkley
- Renee-Coleman-Mitchell
- William Faraclas
- Monika Haugstetter
- Steven Huleatt
- Pamela Kilbey-Fox
- Andrea Lombard
- David Mack
- Patrick McCormack
- Colleen O'Connor
- Elaine O'Keefe
- Joan Segal
- David Skoczulek
- Kristin Sullivan

CPHA Staff

- Eileen Kehl
- Noele Kidney
- Jon Noel

CPHA is an affiliate
of the American Public
Health Association

The Connecticut Public Health Association (CPHA) strongly supports **Senate Bill 16, *An Act Requiring the Labeling of Food and Drink Products That Are Packaged in Materials that Contain Bisphenol-A (BPA)*** as a public health initiative. Labeling of food and beverage packages containing BPA will educate consumers and will enable Connecticut residents to make informed decisions when purchasing food items for their families. CPHA believes that this bill will ultimately benefit the health of Connecticut's citizens by reducing the public's exposure to a chemical of high concern.

BPA is a known endocrine disruptor; a chemical that disrupts the body's endocrine (or hormone) system. BPA is widely used in the manufacturing of polycarbonate plastics and epoxy resins for food and beverage packaging, canned food linings and composite dental fillings and sealants. [1, 2, 14] In fact, nearly one million pounds of products containing BPA are imported or manufactured in the United States each year. [14]

A summary report of the available evidence released by the World Health Organization in 2010 found that use of BPA in food and beverage packaging leads to contamination of the food product it was packaged with. [3, 15] Furthermore, the Centers for Disease Control and Prevention (CDC) have concluded in their latest National Health and Nutrition Examination Survey (2005-2006), that of the 2,638 Americans surveyed 2,548 tested positive for BPA in their urine – these results indicate that 97% of the population is exposed to BPA. [3]

Recent studies have associated exposure to BPA with adverse animal and human health effects. BPA has been associated with prostate and mammary cancer in rodents as well as a multitude of other health risks in these animals. [4, 5] Most importantly, even low doses of BPA in rodents have been shown to cause negative changes in the development of the neurological system and to alter the development of reproductive organs. [5, 12, 7]

The majority of studies on the health effects of BPA have been conducted in animals; however, there is new evidence indicating that there may be an association between exposure to BPA and adverse health conditions in humans. These data show correlations between high levels of BPA in humans and various adverse health effects; however, they cannot determine causality. [17] Experts agree that

more research is needed to determine safe doses of BPA on human health; however, there is growing body of evidence which indicates that the effects of BPA on human health are “complex” and “wide ranging” and are of serious concern. [7, 17, 18]

Infants and young children are thought to be particularly vulnerable to the health effects of BPA. [10, 15] In an effort to prevent harm from exposure to BPA, the U. S. Food and Drug Administration (FDA) released a statement in 2010, recognizing that there is evidence of “potential effects of BPA on the brain, behavior, and prostate gland in fetuses, infants, and young children” and recommended the reduction or replacement of BPA in food can linings as well as baby bottles, cups and formula cans. [9, 10] Moreover, the FDA supports further studies on the evaluation of BPA and is currently working with the World Health Organization, Health Canada and the United Nation’s Agricultural Organization to assess the safety of BPA. [10]

Although U. S. regulatory agencies continue to move slowly regarding regulation of BPA, the Connecticut General Assembly proved its leadership in 2009 when it approved **PA 09-103 AAC Banning Bisphenol A in Children’s Products and Food Products**, and again in 2011, when **PA 11-222 AA Prohibiting the Use of Bisphenol-A in Thermal Receipt Paper** was passed. [10] This trend needs to continue if we are to ensure the safety of our citizens.

The Connecticut Public Health Association supports a complete ban of BPA from food and beverage packaging. Short of this ban, CPHA welcomes **Senate Bill 16 – An Act Requiring the Labeling of Food and Drink Products that are Packaged in Materials that Contain Bisphenol-A** as a step in the right direction toward educating consumers about the presence of BPA in consumer products, with the ultimate goal of reducing exposure. CPHA thanks the Environmental Health Committee in addressing the public health needs of Connecticut’s citizens and appreciates the opportunity to take a stance before the committee today for the labeling of food and beverage packaging containing BPA.

References:

- 1) Vandenberg, L, Hauser, R. Marcus, M., Olea & W. V. Welshons. (2007). Human Exposure to Bisphenol A. *Reproductive Toxicology*, 24(2), 139-177.
- 2) Vandenberg, L., Maffini, M., Schaeberle, C., Ucci, A., Sonnenschein, C., Rubin, B. & A. Soto. (2008). Perinatal Exposure to the Xenoestrogen Bisphenol-A Induces Mammary Intraductal Hyperplasias In Adult CD-1 Mice. *Reproductive Toxicology*, 26(3), 210-219.
- 3) LaKind, J. S. & D. Q. Naimen. (2011). Daily Intake of Bisphenol A and Potential Sources of Exposure: 2005-2006 National Health and Nutrition Examination Survey. *Journal of Exposure Science and Environmental Epidemiology*, 21, 272-279. doi: 10.1038/jes.2010.9
- 4) Jenkins, S., Raghuraman, Eltoum, I., Carpenter, M., et al. (2009). Oral Exposure to Bisphenol A Increases Dimethylbenzanthracene-induced Mammary Cancer In Rats. *Environmental Health Perspectives*, 117(6), 910-915.
- 5) Leranth, C., Hajszan, T., Szigeti-Buck, K., Bober, J. & NJ MacLusky. (2008). Bisphenol A Prevents the Synaptogenic Response To Estradiol In Hippocampus And Prefrontal Cortex Of Ovariectomized Nonhuman Primates. *Proc Natl Acad Sci USA*, 105(37), 14187-14191.
- 6) Lang, I. A. , Galloway, T. & A. Scarlett. (2008). Association of Urinary Bisphenol-A Concentration With Medical Disorders And Laboratory Abnormalities In Adults. *Journal of the American Medical Association*, 300(11), 1303-1310.
- 7) Vom Saal, F.S. et al. (2007). Bisphenol-A Expert Panel Consensus Statement: Integration of Mechanisms, Effects In Animals and Potential To Impact Human Health At Current Levels Of Exposure. *Reproductive Toxicology*, 24(2), 131-138.
- 8) Lee, Y., Ryu, H., Kim. H., Min C., et.al. (2008). Maternal and Fetal Exposure to Bisphenol-A in Korea. *Reproductive Toxicology*, 25, 413-419.
- 9) U. S. Food and Drug Administration. (2012). Bisphenol A (BPA). *U S Food and Drug Administration Home Page*.
- 10) U. S. Food and Drug Administration. (2010 January). Update on Bisphenol A for Use in Food Contact Applications: January 2010. *U S Food and Drug Administration Home Page*. Retrieved from www.fda.gov/NewsEvents/PublicHealthFocus/ucm197739.htm#current
- 11) Connecticut General Assembly. (2012 February 17). 21a-12e Section Text. Retrieved from, search.cga.state.ct.us/dtsearch_pub_statutes.html
- 12) National Toxicology Program Center for The Evaluation of Risks to Human Production. (2008). NTP-CERHR Panel Report on Reproductive and Developmental Toxicity of Bisphenol A. Retrieved from <http://cerhr.niehs.nih.gov/evals/bisphenol/bisphenol.html>
- 13) Shader, M. (2011 June). China Bans BPA From Plastic Baby Bottles. *ConsumerReports.org*. Retrieved from, <http://news.consumerreports.org/safety/2011/06/china-bans-bpas-from-plastic-baby-bottles.html>
- 14) Batra, T. (2011). Bisphenol-A In Canned Food Products: Is It Really Required? *Letter to the Editor*. doi: 10.2478/10004-1254-62-2011-2176
- 15) World Health Organization. (2010). Joint FAO/WHO Expert Meeting to Review Toxicological and Health Aspects of Bisphenol A. Retrieved from, http://www.who.int/foodsafety/chem/chemicals/BPA_Summary2010.pdf
- 16) Kang, J. H., Kondo, F. & Y. Katayama. (2006). Human Exposure To Bisphenol A. *Toxicology*, 226 (2-3), 79-89.
- 17) Rubin, B.S. (2011). Bisphenol A: An endocrine disruptor with widespread exposure and multiple effects. *Journal of Steroid Biochemistry and Molecular Biology*. 127 (1-2), 27-34.
- 18) Keri, R.A.; Ho, S.-M.; Hunt, P.A.; Knudsen, K.E.; Soto, A.M.; Prins, G.S. (2007) An evaluation of evidence for the carcinogenic activity of bisphenol A. *Reproductive Toxicology*. 24(2), 240-252.