

TOXOPLASMOSIS IN PRIMATES

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE ?	TREATMENT	PREVENTION & CONTROL
All nonhuman primate genera infected	Perorally	Fever, anorexia, increased breathing frequency, vomitus, diarrhoea	Yes: in squirrel monkeys and Callitrichidae	Sulfadiazine / Pyrimethamine Spiramycin. Clindamycin	<i>In houses</i> No exposure to feline faeces <i>in zoos</i> no exposure to feline faeces and uncooked meat

Fact sheet compiled by Manfred Brack, formerly German Primate Center, Göttingen / Germany.	Last update November 2003
Fact sheet reviewed by W. Rietschel, Wilhelma Zoologischer-Botanischer Garten, Stuttgart, Germany C. Furley, Howletts Zoo, Bekesbourne, United Kingdom	
Susceptible animal groups All primate genera infected, diseases mostly in New World monkeys and lemurs.	
Causative organism <i>Toxoplasma gondii</i> (Apicomplexa).	
Zoonotic potential Theoretically yes.	
Distribution World- wide.	
Transmission Perorally from cat faeces containing cysts or from meat containing bradyzoites (in man also intrauterine). Aerogenous infection, contamination of bedding material by cat faeces, or uptake of feral rats suspected in captive squirrel monkeys.	
Incubation period Prepatent period after tissue cyst ingestion: 3 – 10 days Prepatent period after ingestion of sporulated cysts : 20 – 34 days.	
Clinical symptoms Fever, anorexia, increased breathing frequency, vomitus, intermittent diarrhoea. In squirrel monkeys frequently death without prior clinical illness.	
Post mortem findings Severe pulmonary edema or focal pneumonia, hydropericard, hydrothorax, focal necroses especially in the liver. In lemurs extensive necrosis of the mesenteric lymph nodes predominated besides necroses in liver and spleen. Cysts in the CNS and other organs, rarely microglial nodules.	
Diagnosis Methylene blue staining, PCR, PAP-immunohistology, direct agglutination, latex agglutination tests, indirect immunofluorescence tests, ELISA, immunosorbent tests. <i>T.gondii</i> infections can be classified using synthetic peptides. Most human infections involve serotypes I to III, but only type I is highly virulent in mice. Cave: <i>Neosporium caninum</i> , which is transmissible at least to Old World monkeys may easily be misdiagnosed as <i>T.gondii</i> in histological slides. confirmation by ELISA or other immunological tests is recommended.	
Material required for laboratory analysis Altered tissues, serum.	
Relevant diagnostic laboratories 1. Local veterinary and medical laboratories. 2. Konsiliarlaboratorium für Toxoplasmose Universitätsklinikum Göttingen Abteilung für Bakteriologie Kreuzberggring 57 D 37075 GÖTTINGEN/Germany	



Tel.: 0551 – 39 – 5801
“ “ 5809
Fax.: “ “ 5861
e-mail: ugross@gwdg.de

Treatment

Sulfadiazine / Pyrimethamine (Daraprim)-combination (In man: 160 mg trimethoprim and 800 mg sulfamethoxazole every other day)
Spiramycin;
Clindamycin hydrochloride (In infants: 300 mg every 6 hs).

Prevention and control in zoos

Prevention of exposure to feline faeces and uncooked meat.. 10 mg /kg benzeneacetonitrile (Diclazuril)

Suggested disinfectant for housing facilities**Notification****Guarantees required under EU Legislation****Guarantees required by EAZA Zoos****Measures required under the Animal Disease Surveillance Plan****Measures required for introducing animals from non-approved sources****Measures to be taken in case of disease outbreak or positive laboratory findings****Conditions for restoring disease-free status after an outbreak****Experts who may be consulted**

Prof.Dr.U.Groß
Koniliarlaboratorium für Toxoplasrose
Univ. Göttingen

References

1. Anderson, D. C., and H. M. McClure. 1993. Toxoplasmosis. *In*: Jones, T. C., U. Mohr, and R. D. Hunt (eds.) Nonhuman Primates II. Monographs on Pathology of Laboratory Animals. Springer, Berlin, Germany. Pp 63-69.
2. Antinori, A., R. Murri, A. Ammassari, A. de Luca, A. Linzalone, A. Cingolani, F. Damiano, G. Maiuro, J. Vacciet, G. Scoppettuolo, E. Tamburrini, and L. Ortona. 1995. Aerosolized pentamidine, cotrimazole and capstone-pyrimethamine for primary prophylaxis of *Pneumocystis carinii* pneumonia and toxoplasmic encephalitis. *AIDS* 9: 1343-1350.
3. Bacciarini, L. N., J. Völlm, M. Janovsky, and A. Gröne. 2001. Acute disseminated toxoplasmosis in a colony of squirrel monkeys (*Saimiri sciureus*) in the Zoologischer Garten Basel, Switzerland. *Verh. Ber. Erkr. Zootiere* 40: 87 – 89.
4. Bowie, W. R., A. S. King, D. H. Werker, J. L. Isaac-Renton, A. Bell, S. B. Eng, and S. A. Marion. 1997. Outbreak of toxoplasmosis associated with municipal drinking water. *Lancet* 350: 173-177.
5. Brack, M. 1987. Agents Transmissible from Simians to Man. Springer, Berlin, Germany.
6. Brack, M., P. Wohlsein, D. Minnemann, and H.-P. Brandt. 1998. Toxoplasmosis outbreak in ringtailed lemurs (*Lemur catta*) and squirrel monkeys (*Saimiri sciureus*). *Primate Rep.* 50: 71-82.
7. Cristina, N., M. L. Dardé, C. Boudin, G. Tavernier, M. Pestre-Alexandre, and P. Ambroise-Thomas. 1995. A DNA-fingerprinting method for individual characterization of *Toxoplasma gondii* strains: Combination with isoenzymatic characters for determination of linkage groups. *Parasitol. Res.* 81: 32-37.
8. Cunningham, A. A., D. Buxton, and K. M. Thomson. 1992. An epidemic of toxoplasmosis in a captive colony of squirrel monkeys (*Saimiri sciureus*). *J. Comp. Pathol.* 107: 207-219.
9. Dietz, H. H., H. Henriksen, V. Bille-Hansen, and S. A. Henriksen. 1997. Toxoplasmosis in a colony of New World monkeys. *Vet. Parasitol.* 68: 299-304.
10. Dreesen, D. W. 1990. *Toxoplasma gondii* infection in wildlife. *J. Am. Vet. Med. Assoc.* 196: 274-276.
11. Dubey, J. P. 1994. Toxoplasmosis. *J. Am. Vet. Med. Assoc.* 205: 1593-1598.
12. Dubey, J. P. 1999. Neosporosis – a newly recognized protozoan disease similar to toxoplasmosis. *Infect. Dis. Rev.* 1: 129 – 130.
13. Epiphanio, S., M. Guimares, D. F. Fedullo, S. H. R. Correa, and J. L. Catos-Dias. 1999. Toxoplasmosis in *Leontopithecus chrysomelas* (Kuhl , 1820) and *Saguinus imperator* (Goeldi , 1907). *Verh. ber. Erkr. Zootiere* 39: 443-444.

14. Escajadillo, A., and J. K. Frenkel. 1991. Experimental toxoplasmosis and vaccine tests in Aotus monkeys. *Am. J Trop. Med. Hyg.* 44: 382-389.
15. Frenkel, J. K. 1990. Transmission of toxoplasmosis and the role of immunity in limiting transmission and illness. *J. Am. Vet. Med. Assoc.* 196: 233-240.
16. Furuta, T., Y. Une, M. Omura, N. Matsutani, Y. Nomura, T. Kikuchi, S. Hattori, and Y. Yoshikawa. 2001. Horizontal transmission of *Toxoplasma gondii* in squirrel monkeys (*Saimiri sciureus*). *Exp. Anim.* 50: 299-306.
17. Gross, U., W. Bohne, M. Soete, and J. F. Dubremetz. 1996. Developmental differentiation between tachyzoites and bradyzoites of *Toxoplasma gondii*. *Parasitol. Today* 12: 30-33.
18. Hall, S. M., A. Pandit, A. Golwilkar, and T. S. Williams. How do Jains get toxoplasma infections? *Lancet* 354: 486-487.
19. Innes, E. A. 1997. Toxoplasmosis: Comparative species susceptibility and host immune response. *Comp. Immunol. Microbiol. Infect. Dis.* 20: 131-138.
20. Juan-Salles, C., N. Prats, A. J. Marco, J. A. Ramos-Vara, D. Borrás, and J. Fernández. 1998. Fatal acute toxoplasmosis in three golden lion tamarins (*Leontopithecus rosalia*). *J. Zoo Wildl. Med.* 29: 55-60.
21. Kong, J. -T., M. E. Grigg, L. Uyetake, S. Parmley, and J. C. Boothroyd. 2003. Serotyping of *Toxoplasma gondii* infections in humans using synthetic peptides. *J. Infect. Dis.* 187: 1484 - 195.
22. Lindsay, D. S., and B. L. Blagburn. 1994. Activity of diclazuril against *Toxoplasma gondii* in cultured cells and mice. *Am. J. Vet. Res.* 55: 530-533.
23. Mazumder, P., H. Y. K. Chuang, M. W. Wentz, and D. L. Wiedbrauk. 1988. Latex agglutination test for detection of antibodies to *Toxoplasma gondii*. *J. Clin. Microbiol.* 26: 244-2446.
24. Pertz, C., R. R. Dubielzig, and D. S. Lindsay. 1997. Fatal *Toxoplasma gondii* infection in golden lion tamarins (*Leontopithecus rosalia rosalia*). *J. Zoo Wildl. Med.* 28: 491-493.
25. Skinner, L. J., J. M. W. Chatterton, A. W. L. Joss, I. L. Moir, and D. O. Ho-Yen. 1989. The use of an IgM immunosorbent agglutination assay to diagnose congenital toxoplasmosis. *J. Med. Microbiol.* 28: 125-128.
26. Une, Y., M. Omura, N. Matsutani, T. Furuta, and Y. Nomura. 1999. Lesions associated with lethal outbreaks of toxoplasmosis in common squirrel monkeys. *Vet. Pathol.* 36: 513.
27. Ven Schoondermark, E. van de, J. Galama, T. Vree, W. Camps, I. Baars, T. Eskes, J. Meuwissen, and W. Melchers. 1995. Study of treatment of congenital *Toxoplasma gondii* infection in rhesus monkeys with pyrimethamine and sulfadiazine. *Antimicrob. Agents Chemother.* 39: 137-144.
28. Wohlsein, P., H.-P. Brandt, M. Brack, M. Peters, G. Schares, und M. Böer. 1999. Immunhistologische und molekularbiologische Untersuchungen von nichthumanen Primaten mit disseminierter Toxoplasmose. *Verh. ber. Erkr. Zootiere.* 39: 209-214.