



MASTOCYTOSIS

<http://www.aocd.org>

Mastocytosis is the term for a group of disorders caused by too many "mast cells". It is most common in children, but the more important forms are seen in both children and adults. Mast cells are found throughout the body, but are especially plentiful in the skin and digestive tract. Mast cells are part of the immune defense system. They are also found around wounds, and seem to play a role in healing of injuries.

Mast cells work by releasing chemicals that attract white blood cells to areas of the body where they are needed. Researchers also think mast cells may have a role in the growth of blood vessels. No one with too few or no mast cells has ever been found, which indicates to some scientists that having too few mast cells may be incompatible with life.

The presence of too many mast cells, or mastocytosis, can occur in two forms: cutaneous and systemic. The cutaneous form is further divided into a diffuse and solitary type. The most common cutaneous (skin) form is also called urticaria pigmentosa (UP), which occurs when mast cells infiltrate the skin.

The solitary form is a single lesion known as a **mastocytoma**.

Systemic mastocytosis is caused by the accumulation of mast cells in the tissues and can affect organs such as the liver, spleen, bone marrow, and small intestine.

Urticaria pigmentosa was first described in 1869, while systemic mastocytosis was first reported in the scientific literature in 1933. The true incidence of either type of mastocytosis remains unknown, but mastocytosis generally is considered to be an "orphan disease." (Orphan diseases affect approximately 200,000 or fewer people in the United States.)

Symptoms

Chemicals released by mast cells cause physiological changes that lead to typical allergic responses such as hives, itching, abdominal cramping, and even shock. Stroking the lesions causes them to itch. When too many mast cells exist in a person's body, the additional chemicals can cause bone pain, abdominal discomfort, nausea and vomiting, ulcers, diarrhea, skin lesions, and episodes of hypotension (very low blood pressure and faintness) or anaphylaxis (shock).

Urticaria pigmentosa can be diagnosed through the abnormally high concentration of mast cells in the skin. Lesions can be erythematous, tan, brown or pink in color.

The diagnosis of systemic mastocytosis can be made by a **biopsy** showing an increased number of mast cells in an organ other than the skin. The biochemical hallmark of the disease is an elevation of histamine or mast cell tryptase protein in blood. Urine may also contain high levels of histamine metabolites (products) as well as metabolites of prostaglandin D2. Elevations of plasma histamine or mast cell protease are occasionally seen in patients with allergic diseases, but unlike mastocytosis patients, their plasma histamine levels are elevated only temporarily. Plasma tryptase and histamine levels are persistently increased in mastocytosis patients.

Other tests that are important in the evaluation of a suspected case of mastocytosis include a bone marrow examination and a bone scan. Special stains, such as Giemsa and toluidine blue, are used on a bone marrow sample to demonstrate the increase in marrow mast cells that occurs in a large percentage of people with the disease.

This information has been provided to you compliments of the American Osteopathic College of Dermatology and your physician.

The medical information provided in this article is for educational purposes only and is the property of the American Osteopathic College of Dermatology. It is not intended nor implied to be a substitute for professional medical advice and shall not create a physician - patient relationship. If you have a specific question or concern about a skin lesion or disease, please consult a dermatologist. Any use, re-creation, dissemination, forwarding or copying of this information is strictly prohibited unless expressed written permission is given by the American Osteopathic College of Dermatology. For detailed information including links to related topics on this and many other skin conditions with photos, visit: <https://www.aocd.org/page/DiseaseDatabaseHome>



MASTOCYTOSIS

<http://www.aocd.org>

Course

Most cases of solitary mastocytoma and UP in children resolve spontaneously. Adults with onset of UP with extensive cutaneous involvement have a higher risk to develop systemic mastocytosis than infants.

Treatment

An array of drugs is used to treat the symptoms of mastocytosis, including **antihistamines** (to prevent mast cell release of chemicals), and anticholinergics (to relieve intestinal cramping). A number of drugs are used for the treatment of specific symptoms of mastocytosis.

This information has been provided to you compliments of the American Osteopathic College of Dermatology and your physician.

The medical information provided in this article is for educational purposes only and is the property of the American Osteopathic College of Dermatology. It is not intended nor implied to be a substitute for professional medical advice and shall not create a physician - patient relationship. If you have a specific question or concern about a skin lesion or disease, please consult a dermatologist. Any use, re-creation, dissemination, forwarding or copying of this information is strictly prohibited unless expressed written permission is given by the American Osteopathic College of Dermatology. For detailed information including links to related topics on this and many other skin conditions with photos, visit: <https://www.aocd.org/page/DiseaseDatabaseHome>