

**STATE ENVIRONMENTAL HEALTH INDICATORS COLLABORATIVE (SEHC)
AIR QUALITY INDICATORS**

Indicator: AQ4: Hazardous Air Pollutants

Measure(s): AQ4.2: Percentage of Population living in counties that exceed health benchmarks for Acrolein

MEASURE DESCRIPTIONS

AQ4.2: Percentage of Population living in counties that exceed health benchmarks for Acrolein	
Last updated :	September 30, 2008
Description:	This analysis compares estimated ambient concentrations of acrolein from EPA's 1999 National Air Toxics Assessment (NATA) with a noncancer health benchmark concentrations. The benchmark concentration corresponds to the level at which exposure to the hazardous air pollutant is judged to be of minimal risk; exposures above this benchmark may be associated with adverse, chronic health effects other than cancer. Note that there are no published cancer health benchmarks for acrolein.
Measurement units:	Percentage of population
Geographic scale:	State summary of County data
Time scale:	1999
Significance/background:	Hazardous air pollutants, also known as air toxics, have been associated with a number of adverse human health effects, including cancers, asthma and other respiratory ailments, and neurological problems such as learning disabilities and hyperactivity. The Clean Air Act identifies 188 substances as hazardous air pollutants. Acrolein is one of the most significant HAPs for noncancer health effects. For detailed background information on the health effects of acrolein, see the following resource: Acrolein - http://www.epa.gov/ncea/iris/subst/0276.htm ; and for information on the setting of HAPs health benchmarks used in this measure, see - http://www.epa.gov/ttn/atw/nata1999/99pdfs/healtheffectsinfo.pdf .
Rationale:	This measure will provide an indication of the population living in counties with levels of acrolein that exceed health benchmark.
Measure limitations:	This measure provides a general indication of the overall population burden of exposure to acrolein. It does not include all of the HAPS. It is based on comparison of estimated ambient concentrations to health benchmarks. Because the database for establishing the health benchmarks is far from complete, this may under-estimate the health significance of these pollutants. The measure does not account for any synergy in effects.
Data resources:	1999 NATA: http://www.epa.gov/ttn/atw/nata1999/tables.html Census: http://www.census.gov/popest/archives/1990s/CO-99-01.html
Data limitations:	The benchmarks generally reflect health risks to adults, rather than potential risks to children or risks in adulthood stemming from childhood exposure. Benchmarks are not available to reflect the latter concerns. Further, the benchmarks reflect risks of continuous exposure over the course of a lifetime. Potential risks from very high short-term exposures are not addressed by these benchmarks.

	The estimates of ambient concentrations of 33 air toxics for the year 1999 were generated as part of EPA's National Air Toxics Assessment. A computer model provided estimates for every county in the continental United States. The computer estimates generally are consistent with the limited set of actual measurements of ambient air toxics concentrations available for 1996, though at many locations the model estimates are lower than the measured concentrations. Actual exposures may differ from ambient concentrations. Indoor concentrations of hazardous air pollutants from outdoor sources may be slightly lower than ambient concentrations, though they can be significantly higher if any indoor sources are present. Levels of some hazardous pollutants may be substantially higher inside cars and school buses.
Related sets of data:	
Related SEHIC measures:	
Related indicators from other projects:	
Recommendations:	
Additional data elements:	

**STATE ENVIRONMENTAL HEALTH INDICATORS COLLABORATIVE (SEHIC)
AIR QUALITY INDICATORS**

Indicator: AQ4: Hazardous Air Pollutants

Measure(s): AQ4.2: Percentage of Population living in counties that exceed health benchmarks for Acrolein

HOW-TO GUIDE: AQ4.2

Obtain Data from the 1999 NATA website:

1. Go to the EPA web site: <http://www.epa.gov/ttn/atw/nata1999/tables.html>
2. In the section **County-Level Ambient Concentration Summaries (Microsoft Access 2000)** select **Acrolein** from the drop-down list of pollutants and then click on the download button.
3. Save the file on your computer, and then open the saved file in Microsoft Excel. To do this, first open Excel, then go under file and choose open, then select **Access Databases** for the Files of Type option. When the Opening Query window pops up, click Open, then select the **County Distributions file** to open.

Create a SEHIC report in Excel:

4. With the county acrolein data now open in Excel, insert a new worksheet in the file by choosing Worksheet under Insert on the drop-down menu at the top of the screen. In the sheet that contains the acrolein data, copy the header row and paste it in the new worksheet. Also highlight all the rows and columns of acrolein data for your state, copy the data, and paste it in the new worksheet.
5. Save the file on your computer as an Excel file.
6. Insert a new column for the noncancer health benchmark after the total_mean column, and label the column noncancer.
 - a. In this new column (noncancer), enter the chronic, noncancer health benchmark of **0.02** in the first cell. Copy this value to the other cells in this column.
7. Now insert a new column and use the IF function to compare the estimated values to the health benchmark, as follows:
 - a. In the new column, enter =IF([column letter and row number corresponding to total_mean value]>[column letter and row number corresponding to the noncancer benchmark value],1,0). (For example, =IF(K4>L4,1,0). K4 and L4 are for illustration; use the column letter and row number that correspond to the indicated data in the formula.) Note: 1=exceeded the benchmark; 0=did not exceed the benchmark.
 - b. Label the new column 'exceeds noncancer'.
8. To calculate the percentage of your state's population in counties where acrolein exceeds the health benchmark:
 - a. Obtain county population data from the census web site <http://www.census.gov/popest/archives/1990s/CO-99-01.html> and copy the 7/1/99 estimates to a new column inserted after the exceeds noncancer column. Label the new column POP.

- b. Insert a new column after the POP column. In the new column, enter the formula =if([column letter and row number of exceeds noncancer]=1,[column letter and row number of population data],0). This formula copies the county population if that county's level of acrolein exceeded the noncancer benchmark. Label this new column POP1.
 - c. Sum the population column to get the state's total population. Sum the pop1 column to get the total population living in counties that exceeded the noncancer benchmark.
9. Open the Excel file SEHIC Indicator Report Template.xls. In the SEHIC Report worksheet AQ4.1, enter these totals in the corresponding cells. Formulas are included in this worksheet for calculating percentages. (These cells will contain #VALUE! until your population counts are entered.)