

State Injury Indicators Report

Instructions for Preparing
2006 Data



Centers for Disease
Control and Prevention
National Center for Injury
Prevention and Control



State Injury Indicators Report: Instructions for Preparing 2006 Data

U.S. Department of Health and Human Services

Centers for Disease Control and Prevention
National Center for Injury Prevention and Control
Division of Injury Response

Atlanta, Georgia
December 2010

State Injury Indicators Report: Instructions for Preparing 2006 Data is a publication of the National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.

Centers for Disease Control and Prevention

Thomas R. Frieden, MD, MPH

Director

National Center for Injury Prevention and Control

Linda Degutis, DrPH, MSN

Director

Division of Injury Response

Richard C. Hunt, MD, FACEP

Director

Suggested citation:

Thomas KE, Johnson RL. State injury indicators report: instructions for preparing 2006 data. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2010.

Editors

Karen E. Thomas, MPH

Division of Injury Response
National Center for Injury Prevention and Control

Renee L. Johnson, RPT, MSPH

Division of Injury Response
National Center for Injury Prevention and Control

Acknowledgements

The editors thank the Safe States Alliance, the Council of State and Territorial Epidemiologists, and their respective members. These partnerships have facilitated the ongoing advancement and success of the development of the injury indicators. The editors also thank Kevin Webb and Bob Thomas, Office of Statistics and Programming, and Angela Marr and Kelly Sarmiento, Division of Injury Response, all with the National Center for Injury Prevention and Control, for their consultation and guidance.

FOREWORD AND UPDATES

The Centers for Disease Control and Prevention's (CDC) National Center for Injury Prevention and Control (NCIPC) is pleased to provide this document to guide you in preparing the 2006 state injury indicators.

Under CDC Program Announcement 05027, 30 states have been funded to collect and submit state injury indicator data; however, all states and U.S. territories are eligible to voluntarily submit data for inclusion in the multistate *State Injury Indicators Report*. As more states and U.S. territories voluntarily participate in this surveillance effort, a broader picture of the burden of injuries can be presented and priorities for prevention can be targeted. During the 2005 data collection cycle, 33 states participated by submitting data for inclusion in the multistate report. We look forward to continuing our work together to advance and improve injury surveillance.

The methods outlined in this document are consistent with those used in previous cycles of injury indicator data collection. These methods are based on recommendations presented in the "Consensus Recommendations for Using Hospital Discharge Data for Injury Surveillance" and in the National Public Health Surveillance System (NPHSS) indicators developed by the Safe States Alliance, (formerly known as the State and Territorial Injury Prevention Directors Association--STIPDA) and the Council of State and Territorial Epidemiologists (CSTE). With partner feedback, CDC continuously modifies and updates the instructions and methodologies outlined in this document.

Changes for the 2006 data collection cycle include:

- In the previous version of the instructions, one of the drowning indicators was inadvertently misnamed. The "Nonfatal Drowning" indicator should have been called "Drowning-Related Hospitalizations" because all hospitalized drownings should be included, even those resulting in in-patient deaths. This has now been corrected.
- Definition pages were added for the Indicators obtained from additional data sources (e.g., Fatality Analysis Reporting System, Behavioral Risk Factor Surveillance System). Because of this, the "Additional Resources" section was revised and moved before the individual indicator pages.

ABBREVIATIONS

BAC	Blood alcohol concentration
BRFSS	Behavioral Risk Factor Surveillance System
CDC	Centers for Disease Control and Prevention
CSTE	Council of State and Territorial Epidemiologists
FARS	Fatality Analysis Reporting System
HDD	Hospital discharge data
ICD-10	International Classification of Diseases – Tenth Revision
ICD-9-CM	International Classification of Diseases – Ninth Revision – Clinical Modification
MVC	Motor vehicle crash
SAVIR	Society for Advancement of Violence and Injury Research
NCCDPHP	National Center for Chronic Disease Prevention and Health Promotion
NCHS	National Center for Health Statistics
NCIPC	National Center for Injury Prevention and Control
NHTSA	National Highway Traffic Safety Administration
NPHSS	National Public Health Surveillance System
STIPDA	State and Territorial Injury Prevention Directors Association
TBI	Traumatic brain injury
VA	Veterans Affairs
WHO	World Health Organization
WISQARS	Web-based Injury Statistics Query and Reporting System
YRBS	Youth Risk Behavior Survey

CONTENTS

Foreword and Updates	iv
Abbreviations	v
Introduction	1
Background and Purpose	3
Preparing the Data Set	5
Additional Resources	9
Injury Indicators	11
All-Injury Indicator 1: Injury Fatalities	12
All-Injury Indicator 2: Hospitalizations for All Injuries	13
Drowning Indicator 1: Unintentional Drowning Fatalities	14
Drowning Indicator 2: Drowning-Related Hospitalizations	15
Fall Indicator 1: Unintentional Fall-Related Fatalities	16
Fall Indicator 2: Unintentional Fall-Related Hospitalizations	17
Fall Indicator 3: Hip Fracture Hospitalizations in Persons Aged 65 Years and Older	18
Fall Indicator 4: Falls in Adults Aged 45 Years or Older	19
Fall Indicator 5: Falls in Adults Aged 45 Years or Older that Caused an Injury	20
Fire-Related Indicator 1: Unintentional Fire-Related Fatalities	21
Fire-Related Indicator 2: Unintentional Fire-Related Hospitalizations	22
Firearm-Related Indicator 1: Firearm-Related Fatalities	23
Firearm-Related Indicator 2: Firearm-Related Hospitalizations	24
Homicide/Assault Indicator 1: Homicides	25
Homicide/Assault Indicator 2: Assault-Related Hospitalizations	26
Motor Vehicle Indicator 1: Motor Vehicle Traffic Fatalities	27
Motor Vehicle Indicator 2: Motor Vehicle Traffic Hospitalizations	28
Motor Vehicle Indicator 3: Seat Belt Use	29
Motor Vehicle Indicator 4: Drinking and Driving	30
Motor Vehicle Indicator 5: Alcohol-Related Crash Deaths	31
Poisoning Indicator 1: Poisoning Fatalities	32
Poisoning Indicator 2: Poisoning Hospitalizations	33
Suicide/Suicide Attempt Indicator 1: Suicides	34
Suicide /Suicide Attempt Indicator 2: Suicide Attempt Hospitalizations	35
Traumatic Brain Injury Indicator 1: Traumatic Brain Injury Fatalities	36
Traumatic Brain Injury Indicator 2: Traumatic Brain Injury Hospitalizations	37
Calculating and Submitting Rates	39
References	41

What is an Injury Indicator?

An injury indicator describes a health outcome of an injury, such as hospitalization or death, or a factor known to be associated with an injury, such as a risk or protective factor among a specified population.

INTRODUCTION

Injury surveillance is one of the most important and basic elements of injury prevention and control. It helps determine the magnitude of injury morbidity and mortality, the leading causes of injury, and the population groups and behaviors associated with the greatest risk of injury. Surveillance data are also fundamental to determining program and prevention priorities. Furthermore, these data are crucial for evaluating the effectiveness of program activities and for identifying problems that need further investigation.

Injury continues to be the leading cause of death and disability among children and young adults.¹ In 2005, more than 173,000 people died from injuries in the U.S. Among them: 26% died from motor-vehicle crashes; 19% died from suicide; and 10% died from homicide.¹ Additionally, in 2005, more than 29 million people were treated for injuries in U.S. emergency departments.¹ The economic cost of injuries is also significant. The total cost of the 50 million medically treated injuries sustained in 2000 is estimated to be \$406 billion in medical expenses and productivity losses.²

The mission of public health includes prevention, mitigation, assurance that the injured have access to treatment, and the reduction of injury-related disability and death.³ The scope of public health encompasses injuries involving any mechanism (e.g., firearm, motor vehicle, or burn) and includes both intentional and unintentional injuries. An important part of the public health mission is to emphasize that injuries are preventable and to dispel the misconception that injuries are unavoidable.

Recognizing the need for more comprehensive injury surveillance data, the State and Territorial Injury Prevention Directors Association (STIPDA) produced *Consensus Recommendations for Injury Surveillance in State Health Departments* in 1999. These recommendations were developed by a working group representing STIPDA; the Council of State and Territorial Epidemiologists (CSTE); the Centers for Disease Control and Prevention (CDC) and its National Center for Injury Prevention and Control (NCIPC); the Society for Advancement of Violence and Injury Research (SAVIR); and individual state partners.

The *State Health Department Consensus Recommendations* identifies specific injuries and injury risk factors to be placed under surveillance by all states and data sets to monitor these injuries and risk factors. The goal is to improve state-based injury surveillance to better support injury prevention programs and policies. By enhancing and standardizing injury surveillance at the state level, its integration with overall public health surveillance as part of the National Public Health Surveillance System (NPHSS) will be much easier.⁴ In tandem with the *State*

Health Department Consensus Recommendations, CSTE and STIPDA developed injury indicators that were formally adopted for inclusion in NPHSS.^{5,6} The NPHSS injury indicators add to other indicators developed by CSTE for chronic diseases and other areas.⁵

The *Consensus Recommendations for Using Hospital Discharge Data for Injury Surveillance*, published in 2003, provides clear and specific recommendations about the evaluation and use of hospital discharge data.⁷ It presents important considerations for the evaluation of data quality and outlines the methodology for developing an injury hospitalization data set.

Collection and dissemination of injury indicators is built upon the foundation laid by the publication of these STIPDA and CSTE documents.

BACKGROUND AND PURPOSE

This manual was created to guide states and U.S. territories in collecting, preparing, and submitting their injury surveillance data. All states and U.S. territories are eligible to voluntarily submit data for this report.

Information obtained from participants will be reviewed and assembled for inclusion in the *State Injury Indicators Report*. This process provides state and U.S. territory injury programs with a standardized method for evaluating injury data and for producing an injury indicator data product that is comparable across states and U.S. territories.

This manual provides straightforward information to encourage participation of all states and U.S. territories regardless of their epidemiologic infrastructure and capabilities. Participation in this report should not be seen as limiting by states of higher capacity, but rather as a place of commonality and a starting point for developing more sophisticated analysis.

The process of preparing indicators is simplified in that it doesn't include the merging and unduplicating of cases found in both hospital discharge and vital statistics data sets. It is important to keep in mind that the quality of the injury indicators is dependant on the completeness and accuracy of external cause coding found on individual state and U.S. territory data sets.

Centralized electronic hospital discharge data and centralized electronic vital statistics data are used to calculate the indicators prepared and submitted by states and U.S. territories. Injuries resulting in or occurring from the following are currently included in the *State Injury Indicators*: all injury, drowning, fall-related injury, fire-related injury, firearm-related injury, homicide/assault, motor vehicle-related injury, poisoning, suicide/suicide attempt, and traumatic brain injury (TBI). Overlap exists among these indicators. For example, a firearm-related homicide would be included in both the firearm-related death indicator and the homicide indicator.

PREPARING THE DATA SET

Background on State Vital Records

Death registration is the responsibility of individual states. The funeral director and the physician who certify the cause of death are usually responsible for the personal and medical information recorded on the death certificate. The cause-of-death section on the certificate is generally the same in all states and is organized according to World Health Organization (WHO) guidelines and coded with ICD-10.⁸ Local registrars assure that deaths in their jurisdictions are registered and that required information is on death certificates before submitting to the state registrar. State registrars number and file the death certificates; certificates of nonresidents are sent to their states of residence. All states send death certificate data to the National Vital Statistics System, managed by CDC's National Center for Health Statistics (NCHS).⁹

Data are limited to information reported on death certificates. The degree of detail in reporting varies among jurisdictions. In general, death certificate data provide limited information about circumstances of injury incidents or contributing factors. The number and type of cause-of-death fields to which states have access also vary, and deaths associated with some injuries, especially suicide, may be underreported. States without access to multiple contributing cause-of-death fields cannot calculate fatality rates for TBI because the diagnostic codes that make up that case definition reside in the contributing cause-of-death fields.

Instructions for Using Vital Statistics Data

Vital statistics data do not require specific preparation for analysis. With the exception of the fatal TBI indicator, all fatal indicators should be calculated by searching the underlying-cause-of-death field only. For the fatal TBI indicator, search *all fields* in the multiple cause of death file. Specific code ranges are identified in the individual indicator specification sheets (see pages 11–42).

Background on State Hospital Discharge Data

At least 90% of all states maintain electronic databases of hospital discharge records for nonfederal, acute care hospitals located within their borders.¹⁰ The information collected varies from state to state. Many states use the standard uniform billing form (UB-92) as the basis for their hospital discharge database. Others use only a subset of variables from the UB-92 for their databases, and a few collect additional variables.

The UB-92, developed by the National Uniform Billing Committee, includes the following data elements:

- patient's age,
- sex,
- zip code,
- admission date,
- length of stay,
- total charges,
- principal diagnosis, and
- up to eight additional diagnoses.

For diagnoses resulting from injuries, an external cause of injury (E-code) is also coded. E-codes, listed in ICD-9-CM, describe several aspects of an injury: intentionality; mechanism; and, for unintentional causes of injury, location of occurrence.¹¹ Completeness of e-coding varies by state.

Instructions for Creating and Using the Injury Hospitalizations Subset of a State Hospital Discharge Data Set

To calculate Injury Hospitalization Indicators, first you need to create an injury subset of hospital discharge records. Create this subset using the following specifications:

- Include only nonfederal, acute care, or inpatient facilities in your hospital discharge data (HDD) injury subset. This excludes Veterans Affairs (VA) and other federal hospitals, rehabilitation centers, and psychiatric hospitals.
- Include readmissions, transfers, and deaths occurring in the hospital.
- Include hospitalizations of state residents only.
- If the data are available, out-of-state hospitalizations of state residents should be included.
- Based on the principal diagnosis field, the subset you create will be injury hospitalizations, defined as follows:
 - Select injury cases by searching only the principal diagnostic code field for the included diagnosis codes. Exclude all other records from the injury hospitalization subset, as shown in the chart below:⁷

INCLUDE	EXCLUDE
800–909.2,	< 800
909.4, 909.9	909.3, 909.5
910–994.9	995.0–995.4
995.5–995.59	995.6–995.7
995.80–995.85	995.86, 995.89
	995.90–995.94
	996–999

Once the injury hospitalization subset has been created, calculate the injury indicators case counts as defined on the individual indicator pages. Search for E-codes in the following manner:


- Search all diagnosis fields.
- If a designated E-code field is in your data set, start with the designated E-code field.
- Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code fields and all diagnostic fields and use the next listed valid E-code. If a case has multiple valid E-codes, then only the first one should be used in the analysis.

- Hospitalizations (except for hip fracture hospitalizations in persons aged 65 years and older) should be age-adjusted to the 2000 standard using the NCHS population distribution (Table 1, page 44).

Assess the completeness and quality measures of the HDD for the following components:

- Percentage of HDD injury records with external cause coding (Figure 1, below).
- Completeness of hospitals participating in the HDD system.
- Inclusion of readmissions and transfers within the data set used for analysis.
- A subjective assessment by health department staff if a substantial proportion of state residents injured in-state are actually hospitalized in a neighboring state.

FIGURE 1.



ADDITIONAL RESOURCES

Other Recommended Data Systems

Indicators based on the Behavioral Risk Factor Surveillance System (BRFSS), the Youth Risk Behavior Survey (YRBS), and the Fatality Analysis Reporting System (FARS) will be calculated at CDC. The data available from YRBS and BRFSS will be examined annually to determine which survey questions should be included.

Behavioral Risk Factor Surveillance System (BRFSS)

CDC's National Center for Chronic Disease Prevention and Health Promotion (NCCDPHP) manages the BRFSS. This is a broad ongoing survey. It is also a state-based, random-digit-dialed telephone survey of the noninstitutionalized U.S. population over age 17. BRFSS monitors risk behaviors associated with the leading causes of disease, injury, and death.¹²

Because BRFSS is telephone-based, population subgroups less likely to have telephones, such as persons of low socioeconomic status, may be underrepresented. In addition, data are self-reported and may be biased. For risk-reduction factors such as self-reported use or testing of smoke alarms, these data may not uniformly represent safe and effective use.¹²

Additionally, not all BRFSS questions are asked every year. Questions asked during the year for which a current Injury Indicator Report is being prepared will be reviewed and appropriate questions included in the report. Results will be reported as a percentage of respondents. For 2006, there are four injury-related BRFSS questions that will be reported.

Youth Risk Behavior Survey (YRBS)

YRBS, a component of the Youth Risk Behavior Surveillance System, is managed by NCCDPHP at CDC. The YRBS monitors risk behaviors associated with the leading causes of injury and death among teenagers.¹³ State and local departments of education and health conduct the survey biennially in many locations throughout the country. The school-based survey is administered to 9th through 12th graders and the data is analyzed by CDC. YRBS data apply only to youth who attend school. The extent of underreporting or overreporting of behaviors cannot be determined, although the survey questions demonstrate good test–retest reliability. Interstate comparisons must be interpreted cautiously because the methods used to collect YRBS data may vary.¹³

Among the 30 funded states, 23 conducted a YRBS in 2005 with overall participation rates of at least 60%.¹⁴ CDC requires a minimum overall participation rate of 60% to generalize the results to the state's population. States with YRBS data meeting this criterion will be included. Results will be reported as a percentage of respondents. No age adjustment will be applied. The YRBS was not administered in 2006.

Fatality Analysis Reporting System (FARS)

FARS, coordinated by the National Highway Traffic Safety Administration (NHTSA), contains data on all fatal traffic crashes that occur in the 50 states, the District of Columbia, and Puerto Rico. For inclusion in FARS, a crash must involve a motor vehicle traveling on a public roadway and result in the death of a person (either a vehicle occupant or a non-motorist) within 30 days of the crash. The FARS file contains a description of each fatal crash reported. More than 100 coded data elements characterize each crash, the vehicles, and the people involved.¹⁵

FARS does not include non-traffic crashes such as those occurring on driveways and other private property. It also does not include deaths occurring more than 30 days after the motor vehicle crash.¹⁵

INJURY INDICATORS

The following pages contain specific case definitions for each of the individual injury indicators. These case definitions should be applied when determining case counts. Once the case counts are determined, they should be entered into the provided spreadsheets for rate calculation and submission to CDC.

ALL-INJURY INDICATOR 1:

Injury Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Injury Fatality ICD-10 Codes

V01–Y36, Y85–Y87, Y89, *U01–*U03	Injury and poisoning
----------------------------------	----------------------

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006—RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Injuries are the leading cause of death for people 1 to 44 years of age and the third leading cause of death overall.¹ Almost 174,000 people died from injuries in 2005.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-13: Reduce deaths caused by unintentional injuries
15-32: Reduce homicides

ALL-INJURY INDICATOR 2: Hospitalizations for All Injuries

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations with any of the following ICD-9-CM diagnostic codes. These should be identified by searching for diagnosis codes only in the principal diagnostic field of the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset). The case count for injury hospitalizations should equal the number of records in your injury hospitalization subset.

Hospitalizations for All Injuries ICD-9-CM Codes

Diagnosis codes

800–909.2, 909.4, 909.9–994.9, 995.5–995.59, 995.80–995.85	Injury and poisoning
---	----------------------

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	Injury is the leading cause of death and disability among children and young adults in the United States. ¹
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding.
HEALTHY PEOPLE OBJECTIVES	15-12: Reduce hospital emergency department visits caused by injuries 15-14: (Developmental) Reduce nonfatal unintentional injuries

DROWNING INDICATOR 1:

Unintentional Drowning Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Unintentional Drowning Fatality ICD-10 Codes

W65–W74	Accidental drowning and submersion
V90	Accident to watercraft causing drowning and submersion
V92	Water-transport-related drowning and submersion without accident to watercraft

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Drowning is one of the 10 leading causes of injury death for persons under age 55 years. In the United States, drowning rates are highest among children under five years of age.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-29: Reduce drownings

DROWNING INDICATOR 2: Drowning-Related Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations with any of the following ICD-9-CM diagnostic or E-codes identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset). These should be identified by searching for diagnosis codes in all diagnostic fields and by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Drowning-Related Hospitalization ICD-9-CM Codes

Diagnosis codes

994.1	Drowning and nonfatal submersion and/or E-codes
E830	Accident to watercraft causing submersion
E832	Other accidental submersion or drowning in water transport accident
E910	Accidental drowning or submersion
E954	Suicide and self-inflicted injury by submersion (drowning)
E964	Assault by submersion (drowning)
E984	Submersion (drowning), undetermined whether accidentally or purposefully inflicted

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population table titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	Drowning-related hospitalizations can result in lifelong disability. Among adolescents and adults, risk factors for drowning include drinking alcohol, swimming alone, and not wearing a personal flotation device while engaged in water sports or recreation. For children under age 5, unexpected access to water or brief lapses in adult supervision are implicated in most drowning incidents. ¹⁷
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.

FALL INDICATOR 1: Unintentional Fall-Related Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Unintentional Fall-Related Fatality ICD-10 Codes

W00–W19	Falls
---------	-------

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006—RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Unintentional falls are the third leading cause of injury death overall and the leading cause of injury death in people 65 years and older.¹ In 2005, there were 19,656 unintentional fall-related deaths.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-27: Reduce deaths from falls

FALL INDICATOR 2: Unintentional Fall-Related Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Unintentional Fall-Related Hospitalization ICD-9-CM Codes

E880–E886, E888	Accidental falls
-----------------	------------------

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006—RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	More than one third of adults 65 and older fall each year. ^{18, 19} Of those who fall, 20% to 30% suffer moderate to severe injuries that make it hard to get around or live alone and increase the chance of early death. ²⁰ The total direct cost of nonfatal fall injuries for people 65 and older in 2000 was \$19 billion. ²¹
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.
HEALTHY PEOPLE OBJECTIVES	No objective

FALL INDICATOR 3:

Hip Fracture Hospitalizations in Persons Aged 65 Years and Older

DEMOGRAPHIC GROUP	Resident persons aged 65 years or older
NUMERATOR	Hospitalizations with the following ICD-9-CM diagnostic code. These should be identified by searching all diagnostic fields of the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset).

Hip Fracture Hospitalization ICD-9-CM Code

Diagnosis code	
820	Fracture of neck of femur

DENOMINATOR	Midyear population of those 65 years and older for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006—RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude. Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	In 2004, there were an estimated 289,000 hospital admissions for hip fractures in people 65 years and older. ²² Up to 25% of adults who lived independently before their hip fracture have to stay in a nursing home for at least a year after their injury ²³ and as many as 20% of hip fracture patients die within a year of their injury. ²⁴
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding.
HEALTHY PEOPLE OBJECTIVES	15-28: Reduce hip fractures among older adults

FALL INDICATOR 4: Falls in Adults Aged 45 Years or Older

This indicator will be calculated at CDC.

DEMOGRAPHIC GROUP	Resident persons aged 45 years or older.
NUMERATOR	Those respondents who experienced a fall.
DENOMINATOR	Respondents aged 45 years or older.
MEASURES OF FREQUENCY	Annual prevalence—crude.
DATA RESOURCES	Data from the Behavioral Risk Factor Surveillance System (BRFSS). ¹²
PERIOD FOR CASE DEFINITION	Past 3 months.
BACKGROUND	More than one third of adults aged 65 years or older fall each year in the United States. ^{18, 19} Many people who fall, even those who are not injured, develop a fear of falling. This fear may cause them to limit their activities, leading to reduced mobility and physical fitness and increasing their actual risk of falling. ²⁵
LIMITATIONS OF INDICATOR	Self-reported data only represent a small portion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of all injuries.
LIMITATIONS OF DATA RESOURCES	As with all self-reported sample surveys, BRFSS data might be subject to systematic error resulting from noncoverage (e.g., lower telephone coverage among populations of low socioeconomic status), nonresponse (e.g., refusal to participate in the survey or to answer specific questions), or measurement (e.g., social desirability or recall bias).
HEALTHY PEOPLE OBJECTIVES	No objective

FALL INDICATOR 5: **Falls in Adults Aged 45 Years or Older that Caused an Injury**

This indicator will be calculated at CDC.

DEMOGRAPHIC GROUP	Resident persons aged 45 years or older.
NUMERATOR	Those respondents who experienced a fall that caused them to limit their regular activities for at least a day or to go see a doctor.
DENOMINATOR	Respondents aged 45 years or older who experienced a fall.
MEASURES OF FREQUENCY	Annual prevalence—crude.
DATA RESOURCES	Data from the Behavioral Risk Factor Surveillance System (BRFSS). ¹²
PERIOD FOR CASE DEFINITION	Past 3 months.
BACKGROUND	More than 3.2 million people aged 45 years or older were treated in emergency departments in 2005 for injuries related to unintentional falls. ¹ This statistic does not include those people who sought care in other settings such as outpatient clinics or doctor's offices.
LIMITATIONS OF INDICATOR	Self-reported data only represent a small portion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of all injuries.
LIMITATIONS OF DATA RESOURCES	As with all self-reported sample surveys, BRFSS data might be subject to systematic error resulting from noncoverage (e.g., lower telephone coverage among populations of low socioeconomic status), nonresponse (e.g., refusal to participate in the survey or to answer specific questions), or measurement (e.g., social desirability or recall bias).
HEALTHY PEOPLE OBJECTIVES	No objective

FIRE-RELATED INDICATOR 1:

Unintentional Fire-Related Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Unintentional Fire-Related Fatality ICD-10 Codes

X00–X09 Exposure to smoke, fire, and flames

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND The United States mortality rate from fires ranks sixth among the 25 developed countries for which statistics are available.²⁶ Four out of five deaths in 2005 occurred in homes²⁷ and approximately half of home fire deaths occurred in homes without fire alarms.²⁸

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-25: Reduce residential fire deaths

FIRE-RELATED INDICATOR 2: Unintentional Fire-Related Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Unintentional Fire-Related Hospitalization ICD-9-CM Codes

E890–E899 Accident caused by fire and flames

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	In 2005, fire departments responded to 396,000 home fires in the U.S., which claimed the lives of 3,030 people (not including firefighters) and injured another 13,825 (not including firefighters). ²⁷ Residential fires disproportionately affect young children, older adults, African Americans, and Native Americans. ²⁹ Working smoke alarms reduce the chance of dying in a house fire by 40% to 50%; however, about 25% of U.S. households lack working smoke alarms. ^{30, 31}
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.
HEALTHY PEOPLE OBJECTIVES	No objective

FIREARM-RELATED INDICATOR 1:

Firearm-Related Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Firearm-Related Fatality ICD-10 Codes

W32–W34	Exposure to inanimate mechanical forces– firearm discharge
X72–X74	Intentional self-harm by firearm discharge
X93–X95	Assault by firearm discharge
Y22–Y24	Firearm discharge of undetermined intent
Y35.0	Legal intervention involving firearm discharge
*U01.4	Terrorism involving firearms

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Firearm-related injuries were the second leading cause of injury-related death in the United States, accounting for about 30,700 deaths in 2005.¹ Nationally, the firearm-related death rate for males is almost seven times higher than that of females.³²

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-3: Reduce firearm-related deaths

FIREARM-RELATED INDICATOR 2: Firearm-Related Hospitalizations

DEMOGRAPHIC GROUP All residents.

NUMERATOR Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949, in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Firearm-Related Hospitalization ICD-9-CM Codes

E922.0–E922.3, E922.8, E922.9	Accident caused by firearm missile
E955.0–E955.4	Suicide and self-inflicted injury by firearms
E965.0–E965.4	Assault by firearms
E985.0–E985.4	Injury by firearms, undetermined whether accidentally, or purposely inflicted
E970	Injury due to legal intervention by firearms
E979.4	Terrorism involving firearms

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006–RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of persons hospitalized. Annual incidence rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Nonfatal firearm-related injury rates are highest among persons ages 15 to 24 years. About one fifth of nonfatal firearm-related injuries treated in U.S. hospital emergency departments are unintentional.³²

LIMITATIONS OF INDICATOR Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.

HEALTHY PEOPLE OBJECTIVES 15-5: Reduce nonfatal firearm-related injuries

HOMICIDE/ASSAULT INDICATOR 1:

Homicides

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Homicide ICD-10 Codes

X85–Y09	Assault
Y87.1	Sequelae of assault
*U01	Terrorism-assault
*U02	Sequelae of terrorism-assault

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Homicide is the fifteenth leading cause of death in the United States; it is the second most common cause of death among persons ages 15 to 24 years.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-32: Reduce homicides

HOMICIDE/ASSAULT INDICATOR 2: Assault-Related Hospitalizations

DEMOGRAPHIC GROUP All residents.

NUMERATOR Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Assault-Related Hospitalization ICD-9-CM Codes

E960–E969	Injury purposely inflicted by other persons
E979	Terrorism
E999.1	Late effect of injury due to terrorism

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006—RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND In 2005, over 1.6 million people were treated in U.S. emergency departments for assault-related injuries with 114,000 of them hospitalized or transferred for a higher level of care.¹

LIMITATIONS OF INDICATOR Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.

HEALTHY PEOPLE OBJECTIVES
 15-34: Reduce the rate of physical assault by current or former intimate partners
 15-37: Reduce physical assaults
 15-38: Reduce physical fighting among adolescents

MOTOR VEHICLE INDICATOR 1:

Motor Vehicle Traffic Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Motor Vehicle Traffic Fatality ICD-10 Codes

V02–V04 (.1, .9), V09.2	Pedestrian injured in transport accident
V12–V14 (.3–.9), V19 (.4–.6)	Pedal cyclist injured in transport accident
V20–V28 (.3–.9), V29 (.4–.9)	Motorcycle rider injured in transport accident
V30–V39 (.4–.9)	Occupant of three-wheeled motor vehicle injured in transport accident
V40–V49 (.4–.9)	Car occupant injured in transport accident
V50–V59 (.4–.9)	Occupant of pick-up truck or van injured in transport accident
V60–V69 (.4–.9)	Occupant of heavy transport vehicle injured in transport accident
V70–V79 (.4–.9)	Bus occupant injured in transport accident
V80 (.3–.5), V81.1, V82.1, V83–V86 (.0–.3), V87 (.0–.8), V89.2	Other land transport accidents

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population table titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Motor vehicle crashes are the leading cause of injury death in the United States. They are also the leading injury cause for years of potential life lost.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that all cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES

- 15-15: Reduce deaths caused by motor vehicle crashes
- 15-16: Reduce pedestrian deaths on public roads
- 26-1: Reduce deaths and injuries caused by alcohol- and drug-related motor vehicle crashes

MOTOR VEHICLE INDICATOR 2: Motor Vehicle Traffic Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Motor Vehicle Traffic Hospitalization ICD-9-CM Codes

E810–E819	Motor vehicle traffic accidents
-----------	---------------------------------

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	In 2005, motor vehicle crashes were the cause of more than 4.3 million emergency department visits in the United States. ¹ It is estimated that front seat occupants who use lap/shoulder belts reduce their risk for fatal injury by about 45% and for moderate to critical injury by 45% to 50%. ³³
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.
HEALTHY PEOPLE OBJECTIVES	15-17: Reduce nonfatal injuries caused by motor vehicle crashes 15-18: Reduce nonfatal pedestrian injuries on public roads 26-1: Reduce deaths and injuries caused by alcohol- and drug-related motor vehicle crashes

MOTOR VEHICLE INDICATOR 3: Seat Belt Use

This indicator will be calculated at CDC.

DEMOGRAPHIC GROUP	Resident persons aged 18 years or older.
NUMERATOR	Those respondents reporting wearing their seatbelt “always” or “almost always” when driving or riding in a car.
DENOMINATOR	Respondents aged 18 years or older.
MEASURES OF FREQUENCY	Annual prevalence—crude.
DATA RESOURCES	Data from the Behavioral Risk Factor Surveillance System (BRFSS). ¹²
PERIOD FOR CASE DEFINITION	No time frame.
BACKGROUND	Safety belts are 45%–60% effective in reducing deaths and 50%–65% effective in reducing moderate-to-critical injuries. ³³
LIMITATIONS OF INDICATOR	Self-reported data only represent a small portion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of all injuries.
LIMITATIONS OF DATA RESOURCES	As with all self-reported sample surveys, BRFSS data might be subject to systematic error resulting from noncoverage (e.g., lower telephone coverage among populations of low socioeconomic status), nonresponse (e.g., refusal to participate in the survey or to answer specific questions), or measurement (e.g., social desirability or recall bias).
HEALTHY PEOPLE OBJECTIVES	15-9: Increase use of safety belts

MOTOR VEHICLE INDICATOR 4: Drinking and Driving

This indicator will be calculated at CDC.

DEMOGRAPHIC GROUP	Resident persons aged 18 years or older reporting drinking at least one alcoholic beverage in the past 30 days.
NUMERATOR	Those respondents reporting driving one or more times after perhaps having too much to drink in the past 30 days.
DENOMINATOR	Respondents aged 18 years or older reporting having a specific number of drinks on one occasion during the previous month (including unknowns and refusals).
MEASURES OF FREQUENCY	Annual prevalence—crude.
DATA RESOURCES	Data from the Behavioral Risk Factor Surveillance System (BRFSS). ¹²
PERIOD FOR CASE DEFINITION	Previous month.
BACKGROUND	In 2005, nearly 1.4 million drivers were arrested for driving under the influence of alcohol or narcotics. ³⁴ This statistic is less than 1% of the 159 million self-reported episodes of alcohol-impaired driving among U.S. adults each year. ³⁵
LIMITATIONS OF INDICATOR	Self-reported data only represent a small portion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of all injuries.
LIMITATIONS OF DATA RESOURCES	As with all self-reported sample surveys, BRFSS data might be subject to systematic error resulting from noncoverage (e.g., lower telephone coverage among populations of low socioeconomic status), nonresponse (e.g., refusal to participate in the survey or to answer specific questions), or measurement (e.g., social desirability or recall bias).
HEALTHY PEOPLE OBJECTIVES	26-1: Reduce deaths and injuries caused by alcohol- and drug-related motor vehicle crashes

MOTOR VEHICLE INDICATOR 5: Alcohol-Related Crash Deaths

This indicator will be calculated at CDC.

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Alcohol-related death of a person involved in crash of a motor vehicle traveling on a public roadway and occurring within 30 days of the crash. Deaths are considered alcohol related if either a driver or nonoccupant (e.g., pedestrian or bicyclist) had a blood alcohol concentration (BAC) greater than or equal to 0.01 g/dL. ¹⁵
DENOMINATOR	Midyear population for the calendar year under surveillance.
MEASURES OF FREQUENCY	Annual number of deaths. Annual mortality rate—crude.
DATA RESOURCES	Fatality Analysis Reporting System (FARS) coordinated by the National Highway Traffic Safety Administration (NHTSA) (numerator) ¹⁵ and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	In 2006, 13,470 people died in alcohol-impaired driving crashes, accounting for nearly one third (32%) of all traffic-related deaths in the United States. Half of the 306 child passengers aged 14 years and younger who died in alcohol-related crashes in 2006 were riding with drivers who had a BAC level of 0.08 g/dL or higher. ³⁶
LIMITATIONS OF INDICATOR	Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less severe injuries.
LIMITATIONS OF DATA RESOURCES	FARS does not include nontraffic crashes such as those occurring on driveways and other private property. In addition, it does not include deaths that occur more than 30 days after the motor vehicle crash. Because blood alcohol levels are not available on all fatalities, the estimates are based on a discriminant analysis of information from all cases where BAC data are available.
HEALTHY PEOPLE OBJECTIVES	15-15: Reduce deaths caused by motor vehicle crashes 15-16: Reduce pedestrian deaths on public roads 26-1: Reduce deaths and injuries caused by alcohol- and drug-related motor vehicle crashes

POISONING INDICATOR 1:

Poisoning Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Poisoning Fatality ICD-10 Codes

X40–X49	Accidental poisoning by and exposure to noxious substances
X60–X69	Intentional self-poisoning
X85–X90	Assault by poisoning
Y10–Y19	Poisoning of undetermined intent
Y35.2	Legal intervention involving gas
*U01 (.6–.7)	Terrorism involving biological or chemical weapons

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Poisoning is the result of the damaging effect of exposure to a broad range of chemicals (e.g., gases, pesticides, heavy metals, drugs, and common household substances such as bleach and ammonia). In 2005, 32,691 people in the United States died from poisoning.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 15-8: Reduce deaths caused by poisonings

POISONING INDICATOR 2: Poisoning Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If a designated E-code field is in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and use the next listed valid E-code.

Poisoning Hospitalization ICD-9-CM Codes

E850–E858	Accidental poisoning by drugs, medicinal substances, and biologicals
E860–E869	Accidental poisonings by other solid and liquid substances, gases, and vapors
E950–E952	Suicide and self-inflicted poisoning
E962	Assault by poisoning
E972	Injury due to legal intervention by gas
E980–E982	Poisoning undetermined whether accidentally or purposefully inflicted
E979 (.6–.7)	Terrorism involving biological or chemical weapons

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	In 1999, 21 states reported that hospitalization rates were 4 to 15 times higher than death rates for poisoning-related injuries. ³⁷
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.
HEALTHY PEOPLE OBJECTIVES	15-7: Reduce nonfatal poisonings

SUICIDE/SUICIDE ATTEMPT INDICATOR 1:

Suicides

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes as an underlying cause of death.

Suicide ICD-10 Codes

X60–X84	Intentional self-harm
Y87.0	Sequelae of intentional self-harm
*U03	Terrorism-intentional self-harm

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND In 2005, suicide was the second leading cause of death among adults ages 25 to 34 years and the third leading cause of death for adolescents and young adults ages 10 to 24 years.¹

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding. The overall completeness of external cause coding on death data is uniformly high. Coding criteria specify that cases of injury death must contain an injury code in the underlying-cause-of-death field.

HEALTHY PEOPLE OBJECTIVES 18-1: Reduce the suicide rate

SUICIDE /SUICIDE ATTEMPT INDICATOR 2: Suicide Attempt Hospitalizations

DEMOGRAPHIC GROUP All residents.

NUMERATOR Hospitalizations identified from the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset) by searching for E-codes in the following manner: Search all diagnosis fields. If there is a designated E-code field in your data set, start with the designated E-code field. Count the first-listed valid E-code, unless it is E849, E967, E869.4, E870–E879, or E930–E949; in which case, search additional E-code and diagnostic fields and then use the next listed valid E-code.

Suicide Attempt Hospitalization ICD-9-CM Codes

E950–E959	Suicide and self-inflicted injury
-----------	-----------------------------------

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND In 2005, there were an estimated 373,000 hospital emergency department visits for suicide attempts in the United States.¹

LIMITATIONS OF INDICATOR Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding. The overall completeness of e-coding is of particular concern and should be reviewed in conjunction with the indicator.

HEALTHY PEOPLE OBJECTIVES 18-2: Reduce the rate of suicide attempts by adolescents

TRAUMATIC BRAIN INJURY INDICATOR 1:

Traumatic Brain Injury Fatalities

DEMOGRAPHIC GROUP All residents.

NUMERATOR Deaths with any of the following ICD-10 codes in any field of the multiple cause of death file.

Traumatic Brain Injury Fatality ICD-10 Codes

S01.0–S01.9	Open wound of head
S02.0, S02.1, S02.3, S02.7–S02.9	Fracture of skull and facial bones
S04.0	Injury of optic nerve and pathways
S06.0–S06.9	Intracranial injury
S07.0, S07.1, S07.8, S07.9	Crushing injury of head
S09.7–S09.9	Other and unspecified injuries of head
T01.0*	Open wounds involving head with neck
T02.0*	Fractures involving head with neck
T04.0*	Crushing injuries involving head with neck
T06.0*	Injuries of brain and cranial nerves with injuries of nerves and spinal cord at neck level
T90.1, T90.2, T90.4, T90.5, T90.8, T90.9	Sequelae of injuries of head

* These codes are not considered valid in the US

DENOMINATOR Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).

MEASURES OF FREQUENCY Annual number of deaths. Annual mortality rate—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population).¹⁶ Rates should be calculated for age and sex.

DATA RESOURCES Death certificate data from vital statistics agencies (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).

PERIOD FOR CASE DEFINITION Calendar year.

BACKGROUND Of the approximately 1.7 million people who sustained a TBI in the United States each year, an estimated 52,000 died; 275,000 were hospitalized; and 1.365 million were treated and released from an emergency department.³⁸

LIMITATIONS OF INDICATOR Injuries severe enough to result in death represent only a small proportion of the overall burden of injury. An evaluation of only these injuries may not present an accurate picture of the causes of less-severe injuries.

LIMITATIONS OF DATA RESOURCES The accuracy of indicators based on codes found in vital statistics data is limited by the completeness and quality of coding.

HEALTHY PEOPLE OBJECTIVES No objective

TRAUMATIC BRAIN INJURY INDICATOR 2: Traumatic Brain Injury Hospitalizations

DEMOGRAPHIC GROUP	All residents.
NUMERATOR	Hospitalizations with any of the following ICD-9-CM diagnostic codes. These should be identified by searching all diagnostic fields of the injury hospital discharge subset (see methods on page 6 for developing the injury hospital discharge subset).

Traumatic Brain Injury Hospitalization ICD-9-CM Codes

Diagnosis codes

800.00–801.99	Fracture of the vault or base of the skull
803.00–804.99	Other and unqualified or multiple fractures of the skull
850.0–850.9	Concussion
851.00–854.19	Intracranial injury, including contusion, laceration, and hemorrhage
950.1–950.3	Injury to the optic chiasm, optic pathways, or visual cortex
959.01	Head injury, unspecified
995.55	Shaken infant syndrome

DENOMINATOR	Midyear population for the calendar year under surveillance. To obtain population estimates by age and sex for your state, use U.S. Census Bureau population tables titled “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT” (see instructions on page 39).
MEASURES OF FREQUENCY	Annual number of persons hospitalized. Annual incidence—crude and age-adjusted (standardized by the direct method to the year 2000 standard U.S. population). ¹⁶ Rates should be calculated for age and sex.
DATA RESOURCES	State hospital discharge data (numerator) and population estimates from the U.S. Census Bureau or suitable alternative (denominator).
PERIOD FOR CASE DEFINITION	Calendar year.
BACKGROUND	An estimated 5.3 million Americans live with a TBI-related disability. According to one study, about 40% of those hospitalized with a TBI had at least one unmet need for services one year after their injury. ^{39, 40}
LIMITATIONS OF INDICATOR	Injuries that result in a hospital admission represent only a portion of the overall burden of injury. Evaluations of these injuries should be considered in the context of both less- and more-severe injuries.
LIMITATIONS OF DATA RESOURCES	The accuracy of indicators based on codes found in hospital discharge data is limited by the completeness and quality of coding.
HEALTHY PEOPLE OBJECTIVES	15-1: Reduce hospitalization for nonfatal head injuries

CALCULATING AND SUBMITTING RATES

Calculation Formula and Instructions

Preformatted rate calculation spreadsheets have been prepared for both the hospital discharge and vital records-based indicators. These spreadsheets can be obtained from Karen Thomas at KThomas@cdc.gov.

Completion of the spreadsheet requires:

- Answering a few data background questions;
- Inserting state population data;
- Entering case counts for individual indicators; and
- Renaming the spreadsheets to reflect state and submission number.

Rate calculations include several types of rates (i.e., age-specific crude rates and age-adjusted rates). The following rate calculation specifications have been preprogrammed into the spreadsheet. If you are preparing these data independent of the spreadsheet, please be sure to follow the same specifications.

- Use the estimated population for the year of the data. This information may be obtained from several sources:
 - www.census.gov/popest/datasets.html (preferred)
 - Scroll to “State population datasets”
 - Continue scrolling to “State Estimates by Demographic Characteristics– Age, Sex, Race, and Hispanic Origin”
 - Continue scrolling to “State Single Year of Age and Sex Population Estimates: April 1, 2000 to July 1, 2006 – RESIDENT”
 - Download File layout
 - Download CSV File
 - your state’s demographic center
- Compute rates per 100,000 population.
- For each indicator, except hip fracture hospitalizations, report age-adjusted rates stratified by sex (female and male), and report the overall age-adjusted rate for the state.
- Report age-specific rates for each indicator in the following age categories:

Under 1

1–4	45–54
5–14	55–64
15–24	65–74
25–34	75–84
35–44	85+

It is possible to obtain the anomalous looking overall age-adjusted rate which does not fall between the two gender-specific age-adjusted rates. Such outcomes are mathematically possible and should be included.

Calculate age-adjusted rates using the age-specific U.S. standard population weights from Table 1.

TABLE 1. AGE ADJUSTMENT TABLE: ALL AGES—ELEVEN AGE GROUPS

Age	U.S. 2000 Standard Population (1,000's)	Adjustment Weights
All ages	274,634	1.000000
Under 1	3,795	0.013818
1–4	15,192	0.055317
5–14	39,977	0.145565
15–24	38,077	0.138646
25–34	37,233	0.135573
35–44	44,659	0.162613
45–54	37,030	0.134834
55–64	23,961	0.087247
65–74	18,136	0.066037
75–84	12,315	0.044842
85+	4,259	0.015508

REFERENCES

1. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. (2005) [cited 2010 Aug 2]. Available from URL: www.cdc.gov/injury/wisqars/index.html.
2. Finkelstein EA, Corso PS, Miller TR, Associates. *Incidence and Economic Burden of Injuries in the United States*. New York: Oxford University Press; 2006.
3. Institute of Medicine (US). *Reducing the Burden of Injury, Advancing Prevention and Treatment*. Washington (DC): National Academy Press; 1999.
4. Meriwether RA. Blueprint for a national public health surveillance system for the 21st century. *J Public Health Manag Pract* 1996;216–23.
5. Council of State and Territorial Epidemiologists. *Injury Control and Prevention Position Statements*. [cited 2010 Aug 2]. Available from URL: <http://www.cste.org/dnn/AnnualConference/PositionStatements/tabid/191/Default.aspx>
6. Safe States Alliance. Injury Surveillance Workgroup. [cited 2010 Aug 2]. Available from URL: <http://www.safestates.org/displaycommon.cfm?an=1&subarticlenbr=10>.
7. Injury surveillance workgroup. *Consensus Recommendations for Using Hospital Discharge Data for Injury Surveillance*. Marietta (GA): State and Territorial Injury Prevention Directors Association; 2003.
8. International Classification of Diseases 10th Revision [online]. [cited 2010 Aug 2]. Available from URL: <http://apps.who.int/classifications/apps/icd/icd10online/>.
9. Centers for Disease Control and Prevention. Mortality Data from the National Vital Statistics System. [cited 2010 Aug 2]. Available from URL: <http://www.cdc.gov/nchs/deaths.htm>.
10. Abellera J, Annet JL, Conn JM, Kohn M. *How states are collecting and using cause of injury data: 2004 update of the 1997 report*. A survey by CSTE, APHA-ICEHS, and STIPDA [online]. [cited 2010 Aug 2]. Available from URL: <http://www.cste.org/pdffiles/newpdffiles/ECodeFinal3705.pdf>.
11. International Classification of Diseases 9th Revision Clinical Modification [online]. [cited 2010 Aug 2]. Available from URL: www.cdc.gov/nchs/icd/icd9cm.htm.
12. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System [online]. [cited 2010 Aug 2]. Available from URL: www.cdc.gov/brfss.
13. Centers for Disease Control and Prevention. Youth Risk Behaviors Surveillance System [online]. [cited 2010 Aug 2]. Available from URL: www.cdc.gov/nccdphp/dash/yrbs/index.htm.
14. Centers for Disease Control and Prevention. Youth Risk Behavior Surveillance—United States, 2005. In: *CDC surveillance summaries*; 2006 Jun 9. *MMWR* 2006;55 (No. SS-5).
15. National Highway Traffic Safety Administration. Fatality Analysis System. [cited 2010 Aug 2]. Available from URL: <http://www.nhtsa.gov/FARS>.
16. Klein RJ, Schoenborn CA. *Age adjustment using the 2000 projected U.S. population*. Hyattsville (MD): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, 2001. Healthy People 2010 statistical notes, No. 20.

17. Fletemeyer JR, Freas SJ, editors. *Drowning: new perspectives on intervention and prevention*. Boca Raton (FL): CRC Press; 1998.
18. Hornbrook MC, Stevens VJ, Wingfield DJ, Hollis JF, Greenlick MR, Ory MG. Preventing falls among community-dwelling older persons: results from a randomized trial. *The Gerontologist* 1994;34(1):16–23.
19. Hausdorff JM, Rios DA, Edelber HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Archives of Physical Medicine and Rehabilitation* 2001;82(8):1050–6.
20. Alexander BH, Rivara FP, Wolf ME. The cost and frequency of hospitalization for fall-related injuries in older adults. *American Journal of Public Health* 1992;82(7):1020–3.
21. Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and nonfatal falls among older adults. *Injury Prevention* 2006;12:290–5.
22. Kozak LJ, DeFrances CJ, Hall MJ. National Hospital Discharge Survey: 2004 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. *Vital Health Stat* 13(162). 2006.
23. Magaziner J, Hawkes W, Hebel JR, Zimmerman SI, Fox KM, Dolan M, et al. Recovery from hip fracture in eight areas of function. *Journal of Gerontology: Medical Sciences* 2000;55A(9):M498–507.
24. Leibson CL, Toteson ANA, Gabriel SE, Ransom JE, Melton JL III. Mortality, disability, and nursing home use for persons with and without hip fracture: a population-based study. *Journal of the American Geriatrics Society* 2002;50:1644–50.
25. Vellas BJ, Wayne SJ, Romero LJ, Baumgartner RN, Garry PJ. Fear of falling and restriction of mobility in elderly fallers. *Age Ageing*. 1997;26:189–93.
26. International Association for the Study of Insurance Economics. *World fire statistics: information bulletin of the world fire statistics*. Geneva (Switzerland): The Geneva Association; 2003.
27. Karter MJ. *Fire loss in the United States during 2005*, Abridged report. Quincy (MA): National Fire Protection Association, Fire Analysis and Research Division; 2006.
28. Ahrens M. *U.S. experience with smoke alarms and other fire alarms*. Quincy (MA): National Fire Protection Association; 2004.
29. Mallonee S, Istre G, Rosenberg M, Reddish-Douglas M, Jordan F, Silverstein P, et al. Surveillance and prevention of residential-fire injuries. *N Eng J Med* 1996;335:27–31.
30. Ahrens M. *U.S. experience with smoke alarms and other fire alarms*. Quincy (MA): National Fire Protection Association; 2001.
31. Smith CL. *Smoke detector operability survey-report findings*. Bethesda (MD): U.S. Consumer Product Safety Commission; 1993 Nov.
32. Gotsch KE, Annet JL, Mercy JA, Ryan GW. Surveillance for fatal and nonfatal firearm-related injuries—United States, 1993–1998. In: *CDC surveillance summaries*; 2001 Apr 13. *MMWR* 2001;50 (No. SS-2).
33. Department of Transportation (US), National Highway traffic Safety Administration. *Traffic safety facts 1999 occupant protection*; 2000; Publication No.: DOT HS 809 090.
34. Crime in the United States 2005: uniform crime reports [database on the Internet]. Washington (DC): US Department of Justice, Federal Bureau of Investigation. 2005 - [cited 2010 Aug 2]. Available from: <http://www.fbi.gov/ucr/05cius/index.html>.

35. Quinlan KP, Brewer RD, Siegel P, Sleet DA, Mokdad AH, Shults RA, Flowers N. Alcohol-impaired driving among U.S. adults, 1993-2002. *Am J Prev Med.* 2005;28(4):345–50.
36. US Department of Transportation, National Highway Traffic Safety Administration (NHTSA). Traffic safety facts 2006: alcohol-impaired driving [monograph on the Internet]. Washington (DC): NHTSA; 2008 [cited 2010 Aug 2]. Available from: <http://www-nrd.nhtsa.dot.gov/Pubs/810801.PDF>.
37. Thomas C, Butler J, Davies M, Johnson R. *State Injury Indicators Report, second edition—1999 data*. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2004.
38. Faul M, Xu L, Wald MM, Coronado VG. *Traumatic Brain Injury in the United States: Emergency Department Visits, Hospitalizations and Deaths 2002–2006*. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2010.
39. Thurman D, Alverson C, Dunn K, Guerrero J, Snieszek J. Traumatic brain injury in the United States: A public health perspective. *J Head Trauma Rehabil* 1999;14(6):602–15.
40. Corrigan JD, Whiteneck G, Mellick D. Perceived needs following traumatic brain injury. *J Head Trauma Rehabil* 2004;19(3):205–16.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Injury Prevention and Control

www.cdc.gov/injury