

PORCINE BRUCELLOSIS (*Brucella suis*)

ANIMAL GROUP AFFECTED	TRANSMISSION	CLINICAL SIGNS	FATAL DISEASE ?	TREATMENT	PREVENTION & CONTROL
Feral and domestic swine (<i>B. suis</i> biovars 1,2,3); European brown hare (<i>B. suis</i> biovar 2); Arctic fox, Wolf, Bear, Reindeer, Moose, Rodents (<i>B. suis</i> biovars 4 and 5)	Spread of <i>B. suis</i> and <i>B. abortus</i> infection in similar way: by artificial insemination or natural mating, occasionally, by arthropods (<i>Oedemagena tarandi</i>)	Generally without any clinical sign; abortion in early and late gestation period can occur. Other symptoms: orchitis, epididymitis, bursitis, synovitis, metritis. General tendency for abscesses to form	No	No	<i>In houses</i> <i>in zoos</i>

<p>Fact sheet compiled by Klaus Gunther Friedrich, Bioparco S.p.A., Rome Zoological Garden, Italy Fabrizio Gamberale, Istituto Zooprofilattico delle Regioni Lazio e Toscana, Rome , Italy</p>	<p>Last update November 2002</p>
<p>Fact sheet reviewed by R. Adone, Istituto Superiore di Sanità, Rome, Italy P. Pasquali, Istituto Superiore di Sanità, Rome, Italy</p>	
<p>Susceptible animal groups Biovars 1, 2 and 3 infect feral and domestic swine. Biovar 2 has been isolated in European brown hare. Biovars 4 and 5 infect, in the arctic region, arctic fox, wolf, bear, reindeer and rodents.</p>	
<p>Causative organism <i>Brucella suis</i>, biovars 1, 2, 3, 4 and 5.</p>	
<p>Zoonotic potential Yes - Biovars 1 and 3 are highly pathogenic for humans. Biovar 2 is rarely pathogenic. Transmission to humans can occur by ingestion, mucosal contact and skin lesions.</p>	
<p>Distribution Biovar 1: South East Asia and Pacific Islands; Latin America. Biovar 2: North East Europe. Biovar 3: North America and China. Biovar 4/5: Arctic Region.</p>	
<p>Transmission <i>B. suis</i> infection is transmitted by mucosal contact with contaminated materials such as faeces or uterine discharges and vaginal fluids. Transmission can also occur by venereal route or, occasionally, by arthropods (<i>Oedemagena tarandi</i>).</p>	
<p>Incubation period Variable.</p>	
<p>Clinical symptoms Generally without any clinical sign, or orchitis, epididymitis, bursitis, synovitis, metritis, abortion.</p>	
<p>Post mortem findings Depending on <i>B. suis</i> biovar involved. The most common findings are: nodular and miliar lesions in the uterus,</p>	



purulent metritis, endothelial sclerosis of uterine vessels, orchitis, epididymitis, arthritis, spondylitis, granulomatous lesions in skeleton and joints, vertebral osteomyelitis, abscesses.

Diagnosis tests

- Direct diagnosis: Polymerase chain reaction (PCR), Culture for isolation.
- Indirect diagnosis: Complement Fixation Test (CFT), Milk Ring Test, ELISA-s, Brucellin skin test

Material required for laboratory analysis

For indirect diagnosis: blood serum and/or milk.

For direct diagnosis: blood, vaginal swabs, foetal membranes, materials from aborted foetus such as spleen, stomach contents, encephalon and lung.

EU Reference Laboratory**AFSSA, Nancy**

Laboratoire d'études sur la rage et la pathologie des animaux sauvages

Domaine de Pixérécourt, BP 9

F-54220 Malzéville

France

OIE Reference Laboratories

- **Dr Heinrich Neubauer**
Federal Research Centre for Virus Diseases of Animals (BFAV), Institute of Bacterial Infections and Zoonoses
Naumburger Str. 96a, 07743 Jena
GERMANY
Tel: (49.3641) 80.42.00 Fax: (49.3641) 80.42.28
Email: heinrich.neubauer@fli.bund.de
- **Dr Ana Maria Nicola**
Gerencia de Laboratorios (GELAB), Servicio Nacional de Sanidad y Calidad Agroalimentaria (SENASA)
Av. Alexander Fleming 1653, 1640 Martínez, Pcia. de Buenos Aires
ARGENTINA
Tel: (54.11) 48.36.19.92 Fax: (54.11) 48.36.19.92
Email: anicola@senasa.gov.ar
- **Dr J.A. Stack**
VLA Weybridge
New Haw, Addlestone, Surrey KT15 3NB
UNITED KINGDOM
Tel: (44.1932) 35.76.10. Fax: (44.1932) 35.72.16
Email: j.a.stack@vla.defra.gsi.gov.uk
- **Dr B. Garin-Bastuji**
AFSSA Alfort, Unité Zoonoses Bactériennes, Lab. OIE/FAO de référence pour la brucellose animale, Laboratoire d'études et de recherches en pathologie animale et zoonoses
23 avenue du Général de Gaulle, 94706 Maisons-Alfort Cedex
FRANCE
Tel: (33 (0)1) 49.77.13.00 Fax: (33 (0)1) 49.77.13.44
Email: b.garin-bastuji@afssa.fr
- **Dr K. Nielsen**
Canadian Food Inspection Agency, Animal Diseases Research Institute
P.O. Box 11300, Station H, Nepean, Ontario K2H 8P9
CANADA
Tel: (1.613) 228.66.98 ext. 48.04 Fax: (1.613) 228.66.69
Email: nielsenk@inspection.gc.ca
- **Dr Massimo Scacchia**
CESME, Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise 'G. Caporale', National Centre for Exotic Diseases
Via Campo Boario, 64100 Teramo
ITALY



Tel: (390.861) 33.24.05 Fax: (390.861) 33.22.51
Email: m.scacchia@izs.it

Treatment

No treatment.

Prevention and control in zoos

See fact sheet for *B. abortus*.

Suggested disinfectant for housing facilities

See fact sheet for *B. abortus*.

Notification

Yes.

Guarantees required under EU Legislation

- Directive 92/65/EEC
- Directive 64/432/EEC
- Directive 97/12/EEC
- Directive 97/12/EEC
- Directive 64/432/EEC
- Directive 92/65/EEC
- Directive 90/425/EEC
- Directive 89/662/EEC
- Directive 1999/466/EEC
- Directive 1999/467/EEC

Guarantees required by EAZA Zoos**Measures required under the Animal Disease Surveillance Plan****Measures required for introducing animals from non-approved sources****Measures to be taken in case of disease outbreak or positive laboratory findings****Conditions for restoring disease-free status after an outbreak****Contacts for further information**

Dr. Fabrizio Gamberane, Email fgamberale@rm.izs.it

References

1. Alton, G. G., L. M. Jones, R. D. Angus, and J. M. Verger. 1988. Techniques for the brucellosis laboratory. I.N.R.A., Paris, France.
2. Thorne, E. T. 2001. Brucellosis. *In*: Infectious diseases of wild mammals, E. S. Williams and I.K. Barker (eds). Manson Publishing, London, UK. Pp. 372-395.